

Dr. David Kirlin, DPM

Dr. Ryan Meredith, DPM

Dr. Wagner Santiago, DPM



WART SURGERY INSTRUCTIONS

GENERAL INFORMATION

Stay off your feet as much as possible today. You may wear any shoe that does not squeeze, constrict or put pressure on the surgical area. Your foot may remain numb for up to 6–10 hours after the procedure. Please contact our office if you have any questions or concerns.

Post–Op Instructions

1. Remove bandage the day following your surgery.
2. Apply AmeriGel or a triple antibiotic ointment sparingly to wound, then apply a dry dressing and paper tape. Do this once a day for 7 days or until healed once the initial bandage has been removed.

Bleeding– Slight bleeding, discoloration and drainage are normal. If bleeding comes through the dressing, elevate and reinforce the dressing with extra gauze.

Discomfort– You can elevate your foot to help alleviate minor swelling, bleeding and discomfort. You may also take Tylenol or other over–the–counter pain relievers as directed on the package. If the pain or discomfort is not controlled adequately, then contact our office.

Removing the surgical bandage– The day after your surgery, carefully remove the dressing and shower or bathe as normal. If the gauze or bandage sticks to the area, dampen it with water or shower/bathe with the bandage in place. This will make it easier to remove with minimal or no discomfort.

DRESSING SHOULD BE CHANGED ONCE DAILY FOLLOWING BATHING